

Antioxidant activity and physicochemical stability of phytocosmetic formulations containing *Phyllanthus niruri* extract

Heitor Janaudis Carmagnani¹ , Gabriel Bucciarelli Mansano¹ , Priscila Alves Balista¹ , Rodrigo Vieira Gonzaga^{1,2} , Flávia Sobreira¹ , Michelli Ferrera Dario^{1,2,*} 

¹Curso de Farmácia, Centro Universitário São Camilo, 04263-200, São Paulo, Brazil

²Departamento de Farmácia, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, 05508-000, São Paulo, Brazil

*Corresponding author: mfdario@gmail.com

ABSTRACT

Antioxidants are an appreciable strategy to prevent skin photoaging as they inhibit reactive oxygen species formation under UV exposure. *Phyllanthus niruri* L. (Euphorbiaceae) is a medicinal plant utilized in treating urolithiasis and contains high amounts of secondary metabolites with antioxidant activity. The potential cosmetic usage of the dry extract of *P. niruri* in three cosmetic base formulations (non-ionic emulsion, anionic gel, and gel-cream) were assessed by evaluating antioxidant activity and thermal stability. Antioxidant activity was assessed by the DPPH method and the preliminary stability test was performed in different storage temperature

conditions (40.0±2.0, 20.0±5.0, 5.0±2.0 °C). Organoleptic (appearance, color, odor), sensory (consistency) and physical-chemical characteristics (pH) were evaluated after 30 days. Formulations containing the plant extract presented high antioxidant activity, whereas the non-ionic emulsion had the best performance. All formulations exhibited antioxidant activity stability when stored at low temperatures, however, medium and high temperatures caused a reduction of this parameter. Therefore, non-ionic emulsion was considered the ideal base formulation for antiaging purposes as long as stored at low temperature.

Keywords: Free radical scavenger, Phenolic compound, Semi-Solid Formulation, Cosmetic.

INTRODUCTION

Skin aging is a natural and complex biological process characterized by a reduction in the skin's antioxidant capacity over time. However, extrinsic factors, such as chronic exposure to ultraviolet (UV) radiation, are responsible for photoaging, a type of premature aging characterized by deep wrinkles, hyperpigmentation or hypopigmentation, reduced elasticity, and drier and rougher skin. UV radiation stimulates the production of reactive oxygen species (ROS) in the dermis, which may damage cell membranes, proteins, enzymes, and even DNA through a chain oxidation reaction that can induce cell death and skin cancer (Shanbhag et al. 2019).

Photoprotection and reducing sun exposure are well-accepted strategies to prevent premature aging (Kostyuk et al. 2018). Furthermore, antioxidant use helps prevent aging as it captures ROS and inhibits

cell damage. As such, antioxidant phytochemicals are interesting active pharmaceutical ingredients in skincare products (Oliveira and Dario 2019).

Phyllanthus niruri L. (Euphorbiaceae) is a medicinal plant found in tropical and subtropical regions, especially Brazil, India, Africa, Malaysia, and Indonesia (Kumar et al. 2017). The species has several pharmacological properties, including antibacterial, anti-inflammatory, hepatoprotective, antiulcer, and antidiarrheal properties; however, it is mainly used in treating urolithiasis. These properties are due to its secondary metabolites, such as tannins, flavonoids, alkaloids, terpenes, coumarins, and lignans (Lee et al. 2016; Kaur et al. 2017). *P. emblica* L., known as emblica or amla, is a species of the same genus that has demonstrated antioxidant capacity as it has inhibited metalloproteinases related to skin aging, protecting fibroblasts from

Received: March 27, 2023

Accepted after revision: August 31, 2023

Published on line: September 30, 2023

ISSN 1983-084X

© 2023 Revista Brasileira de Plantas Medicinais/Brazilian Journal of Medicinal Plants.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

oxidative stress and stimulating their proliferation as well as increasing the production of Type I collagen (Chaikul et al. 2021).

Moreover, as most *P. niruri* compounds have antioxidant activity, this study aimed to evaluate this extract as a candidate for antiaging products.

MATERIAL AND METHODS

Chemicals

Disodium EDTA; a mixture of cetearyl alcohol, polysorbate 60, PEG-150 stearate, and steareth-20 (Polawax® NF); dimethicone; butylated hydroxytoluene; ethylhexyl stearate; C10-30 alkyl acrylate crosspolymer (Carbopol® 920); hydroxyethylcellulose; glycerin; methylparaben; propylparaben; aminomethyl propanol (AMP® 95); propylene glycol; and caprylic capric triglyceride were used without further purification. The solvents ethanol (Prolink) and methanol (Anidrol) were PA grade. Gallic acid (purity > 98.0%) and 2,2-diphenyl-1-picrylhydrazyl (DPPH) were purchased from Sigma Aldrich. *P. niruri* was purchased from SantosFlora Comércio de Ervas Ltda., corresponding to Lot 1608175268. This research was recorded (A45E2DF) in the National System for the Management of Genetic Heritage and Associated Traditional Knowledge (SisGen).

Phyllanthus niruri extract

The dried *P. niruri* aerial plant was pulverized in a knife mill and sieved in 20 mesh sieves (850 µm). Extraction was performed through ultrasound at 22 °C for 30 min using a proportion drug to extractor liquid (ethanol 50% v/v) ratio of 1:10 (mass of the plant drug versus liquid extractor). Subsequently, the ethanol was removed in a rotary evaporator under reduced pressure at 40 °C. The remaining aqueous extract was dried and stored in a desiccator at 22 °C. Thin layer chromatography profile of dried *P. niruri* extract was performed by our group as shown at Carmagnani et al. (2020).

Preparation of cosmetic formulations

The dried *P. niruri* extract was incorporated at 2.5% (w/w) into three cosmetic bases: non-ionic emulsion, anionic gel, and gel-cream (Table 1). The percentage of flavonoids in the extract, previously shown at Carmagnani et al. (2020), was used to define the concentration of the extract in formulations.

Preliminary stability test

Formulations were subjected to a preliminary stability test to evaluate the occurrence of physical and organoleptic characteristics (color,

Table 1. Composition of base cosmetic formulations.

Ingredients	INCI*	Non-ionic emulsion	Anionic gel	Gel- cream
		% (w/w)		
EDTA	Disodium EDTA	0,1	0,1	0,1
-	-	-	-	-
Deionized water	Aqua	qsp 100	qsp 100	qsp 100
Polawax® NF	Cetearyl Alcohol (and) Polysorbate 60 (and) PEG-150 Stearate (and) Steareth-20	15	-	15
Dimethicone	Dimethicone	2	-	-
Butylated hydroxytoluene (BHT)	BHT	0,05	-	0,2
Ethylhexyl Stearate	Ethylhexyl Stearate	2	-	-
Carbopol® 920	C10-30 Alkyl Acrylate Crosspolymer	-	0,8	-
Hydroxyethylcellulose	Hydroxyethylcellulose	-	-	0,5
Glycerin	Glycerin	-	5	-
Methylparaben	Methylparaben	0,2	0,15	0,2
Propylparaben	Propylparaben	0,1	-	0,2
AMP 95	Aminomethyl Propanol	-	qs pH 6,5-7,0	-
Propylene Glycol	Propylene Glycol	3,0	-	8
Caprylic Capric Triglyceride	Caprylic Capric Triglyceride	-	-	10

odor, and aspect), sensory, pH value alterations, and antioxidant activity. The formulations were subjected to different storage temperature conditions (40.0 ± 2.0 , 5.0 ± 2.0 , and 20.0 ± 5.0 °C) for 30 days. Furthermore, pH values were measured through the direct immersion of an electrode in the formulations using a potentiometer, and antioxidant activity was determined using a DPPH assay.

Evaluation of antioxidant activity

Antioxidant activity of all formulations, including blank (no added extract), was assessed using the DPPH method. First, 0.2 g of each formulation was added to 5 ml of methanol PA and vortexed. Subsequently, the formulations were centrifuged at $600 \times g$ for 15 min. About 2.5 ml of the supernatant was transferred to a 10 ml volumetric flask and its volume was completed with ethanol PA. An aliquot of 500 µl of these solutions was added to 2.5 ml of 100 µM DPPH ethanolic solution. The reaction proceeded for 30 min in the dark, and the spectrophotometric reading was performed at 517 nm using absolute ethanol as a reading blank. Negative control was performed by adding 0.5 ml of ethanol PA and 2.5 ml of DPPH solution. The solutions were prepared in triplicate (Dario et al. 2013). The percentage of free radical scavenging (% FRS) was calculated following Equation 1.

$$\% \text{ FRS} = \frac{Abs_C - Abs_S}{Abs_C} \quad (\text{Eq. 1})$$

where: Abs_C : Negative control absorbance; Abs_S : Sample absorbance.

The antioxidant activity was expressed in gallic acid equivalents (GAE). A calibration curve was constructed from serial dilutions of reference standard gallic acid (1.5 - 12.0 µg/ml) in methanol PA. An aliquot of 500 µl of these solutions was added to 2.5 ml of 100 µM DPPH ethanolic solution, and the protocol proceeded as described above. The antioxidant capacity was determined by comparing the % FRS of each formulation to the gallic acid calibration curve and reported in mg GAE/g formulation.

Statistical analysis

Statistically significant differences in antioxidant activity during preliminary stability test were verified using a paired *t*-test comparing initial and final values (after 30 days of stability test) ($\alpha = 0.05$).

RESULTS AND DISCUSSION

Formulations based on emulsions, gels, and gel-creams are the most common vehicles employed in cosmetics due to their excellent sensorial characteristics and acceptance by consumers (Soares et al. 2021). Emulsions are composed of two immiscible phases (aqueous and oily) stabilized by surfactants whose charge (anionic, cationic, non-ionic or amphoteric) determines the type of emulsion. This type of formulation has moisturizing and emollient properties. Gels are semisolid dosage forms, usually water-based, thickened by a gelling agent. They are cosmetically elegant, nongreasy and provide a cooling sensation (Mayba and Gooderham 2018). By the other hand, gel-creams are emulsions thickened by a hydrophilic colloidal consistency agent (Ferreira et al. 2018).

























The organoleptic and sensory characteristics of a product determine its acceptance by the consumer as the analysis of these characteristics guarantees that any changes in the product do not alter its efficacy, safety or acceptability. Evaluating these characteristics provides parameters to estimate the state of the formulation under study through comparative analysis (i.e., the state of the formulation before and after going through certain processes). Parameters such as appearance, color, and odor are usually evaluated macroscopically (Rosário et al. 2021).

In general, the preliminary stability test showed no profound alterations in the formulations (Table 2). One of the alterations was the milder odor of the anionic gel containing the plant extract when stored at a high temperature (40.0 ± 2.0 °C). This may be due to the evaporation of ethanol from the sample or various reactions, such as oxidation, hydrolysis, or interactions with the packaging material, that can reduce the number of substances in the extract (Casteli et al. 2008).

The non-ionic emulsion changed from a viscous to a firmer appearance, possibly owing to the loss of water from the formulation through evaporation (Pianovski et al. 2008). When opening the pots, small droplets of water were seen on the lids. This phenomenon was probably caused by the insufficient concentration of humectants, which can prevent water loss (Souza and Ferreira 2010). The addition of glycerin, for example, could solve instability.

The pH of the formulation is an important parameter because it evidences alterations and reactions that can modify the pH of the formulation. The pH of healthy skin is slightly acidic (4.5–6.0). This acidity is important for maintaining the homeostasis of the skin's barrier function (physical and chemical protection) and protecting against microorganisms, especially those alien to the human microbiota

Table 2. Aspect, color and pH values of formulations subjected to preliminary stability test.

Formulation		20.0±5.0 °C	40.0±2.0 °C	5.0±2.0 °C	
Anionic gel base					
pH		5,78	5,1	6,2	5,18
Anionic gel + <i>P. niruri</i> extract					
pH		6,28	6,52	5,46	6,35
Non-ionic emulsion base					
pH		5,59	4,31	4,62	5,74
Non-ionic emulsion + <i>P. niruri</i> extract					
pH		7,2	5,03	5,62	6,6
Gel-cream base					
pH		5,57	6,99	4,86	4,41
Gel-cream + <i>P. niruri</i> extract					
pH		5,93	5,22	5,27	5,26




Figure 1. Antioxidant activity of the formulations during preliminary stability test. The formulations that contain an asterisk (*) have statistically different antioxidant activity ($p \leq 0.05$) compared to the initial time (t_0). The values are expressed by the average antioxidant activity (triplicate) and error bars represent the standard deviation.

(such as bacteria and fungi) (Leonardi et al. 2002). To maintain healthy skin, topical formulations must maintain an acidic pH (~5 is desirable). Formulations with $pH < 5$ help maintain skin homeostasis, thus preventing aggressive agents, such as microorganisms, as well as dermatitis, acne, and eczema (indicating unhealthy skin) (Das and Wong 2020).

The formulations in our study showed changes in pH as they were stored at different temperatures (Table 2). Monitoring pH in cosmetic formulations are essential as organoleptic characteristics (appearance, color, and odor) and the effectiveness of the active ingredient can be changed as a result of the pH changes. Additionally, phytocosmetics containing polyphenols (tannins, coumarins, and flavonoids) have increased stability at a slightly acidic pH (Anvisa 2021).

Carbomer gels, such as Carbopol®, are stable at a pH of 5.5–7.3, presenting the ideal viscosity. The gel-cream and non-ionic emulsion must have a pH of 5.5–6.5. Therefore, the gel remained at an ideal pH during the preliminary stability test, whereas the non-ionic emulsion and gel-cream showed a pH (4.31–6.99) outside the established ideal range but within the pH range biocompatible with the skin (Aulton 2005).

The antioxidant activity of the formulations and extract was expressed in gallic acid equivalents based on a calibration curve obtained for this primary reference standard (Figure 1). The base formulations (gel-cream, non-ionic emulsion, and anionic gel) presented extremely low antioxidant activity as

the dry *P. niruri* extract was not incorporated. The antioxidant activity of the gel-cream base was higher than the other formulations, possibly due to its increased concentration of BHT, which is an antioxidant. Formulations containing the dry extract showed considerable antioxidant activity. Owabhel and Eboh (2022) demonstrated that DPPH scavenging of *Phyllanthus niruri* methanolic extract was slightly smaller than gallic acid, but the nitric oxide scavenging was higher. This result can be explained by the presence of phenolics, such as rutin, quercetin, niruriflavone, and other potent antioxidants phytochemicals (Bagalkotkar et al. 2006, Ferrante et al. 2020). The addition of the *P. niruri* extract caused an increase of 936, 2863, and 2161% in gel-cream, anionic gel and non-ionic emulsion, respectively. The non-ionic emulsion showed superior values of antioxidant activity before the stability test (t_0).

No statistical difference was observed for formulations containing the extract when stored at low temperatures ($5.0 \pm 1.0^\circ \text{C}$), while all formulations stored in an oven ($40.0 \pm 2.0^\circ \text{C}$) and at room temperature ($20.0 \pm 5.0^\circ \text{C}$) showed significant differences. These data indicate that storage at increased temperatures can cause alterations in the formulation, such as the formation of degradation products and loss of activity of the active pharmaceutical ingredient; therefore, low-temperature storage is recommended (Lange et al. 2009).

Temperature directly influences the conservation of plant drugs. In general, the plants

must remain at room temperature (less than 15 and up to 30 °C) without the incidence of light and protected from humidity. High temperatures (above 50 °C) can degrade plant compounds, causing the loss of pharmacological activity through the formation of other compounds (Oliveira 2014).

Our findings show that *P. niruri* extract, rarely used for cosmetics, has excellent potential for antiaging products due to its high antioxidant activity. The base cosmetic formulation directly influenced the physical and chemical stability of the formulations, and non-ionic emulsion was most suitable for incorporating the dry *P. niruri* extract since it is stored at low temperature conditions.

ACKNOWLEDGEMENT

We thank Centro Universitário São Camilo and Profa. Dra. Maria Valéria Robles Velasco (Faculty of Pharmaceutical Sciences - University of São Paulo) for the research support.

AUTHORS' CONTRIBUTION

Conceptualization, F.S.M.G. and M.F.D.; methodology, M.F.D.; validation, F.S.M.G. and M.F.D.; formal analysis, M.F.D.; investigation, H.J.C. and G.B.M.; resources, F.S.M.G. and M.F.D.; data curation, M.F.D.; Writing - original draft, H.J.C., G.B.M. and R.V.G.; Writing - review & editing, M.F.D. and P.A.B.; visualization, R.V.G.; supervision, M.F.D.; project administration, M.F.D.; funding acquisition, M.F.D. All authors read and approved the final manuscript.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

REFERENCES

Anvisa (2021) Formulário de Fitoterápicos da Farmacopeia Brasileira. 2.ed. Brasília: Anvisa. 216p.

Aulton M (2005) Delineamento de formas farmacêuticas. 2.ed. Porto Alegre: Artmed. 678p.

Bagalkotkar G, Sagineedu SR, Saad MS, Stanslas J (2006) Phytochemicals from *Phyllanthus niruri* Linn. and their pharmacological properties: a review. J Pharm Pharmacol 58:1559-1570.

Carmagnani HJ, Mansano GB, Sobreira F (2020) Otimização do processo extrativo de *Phyllanthus niruri* L. Mundo saúde 44:134-143. <https://doi.org/10.15343/0104-7809.202044134143>

Casteli VC, Mendonça CC, Campos MAL, Ferrari M, Machado SRP (2008) Desenvolvimento e estudos de estabilidade preliminares de emulsões O/A contendo cetoconazol 2,0%. Acta Sci Health Sci 30:121-128. <https://doi.org/10.4025/actascihealthsci.v30i2.812>

Chaikul P, Kanlayavattanukul M, Somkumnerd J, Lourith

N (2021) *Phyllanthus emblica* L. (amla) branch: A safe and effective ingredient against skin aging. J Tradit Complement Med 11:390-399. <https://doi.org/10.1016/j.jtcm.2021.02.004>

Dario MF, Pahl R, Castro JR, Lima FS, Kaneko TM, Pinto CASO, Baby AR, Velasco MVR (2013) Efficacy of *Punica granatum* L. hydroalcoholic extract on properties of dyed hair exposed to UVA radiation. J Photochem Photobiol B 120:142-147. <https://doi.org/10.1016/j.jphotobiol.2012.12.011>

Das S, Wong ABH (2020) Stabilization of ferulic acid in topical gel formulation via nanoencapsulation and pH optimization. Sci Rep 10:12288. <https://doi.org/10.1038/s41598-020-68732-6>

Ferrante C, Chiavaroli A, Angelini P, Venanzoni R, Flores GA, Brunetti L, Petrucci M, Politi M, Menghini L, Leone S, Recinella L, Zengin G, Ak G, Mascio M, Bacchin F, Orlando G (2020) Phenolic content and antimicrobial and anti-inflammatory effects of *Solidago virga-aurea*, *Phyllanthus niruri*, *Epilobium angustifolium*, *Peumus boldus*, and *Ononis spinosa* extracts. Antibiotics 9:783. <https://doi.org/10.3390/antibiotics9110783>

Ferreira AO, Brandão M, Polonini HC (2018). Guia Prático da Farmácia Magistral. 5.ed. São Paulo: Pharmabooks. 1690p.

Kaur N, Kaur B, Sirhindi G (2017) Phytochemistry and pharmacology of *Phyllanthus niruri* L.: A review. Phytother Res 31:980-1004. <https://doi.org/10.1002/ptr.5825>

Kostyuk V, Potapovich A, Albuhaydar AR, Mayer W, Luca C, Korkina L (2018) Natural substances for prevention of skin photoaging: screening systems in the development of sunscreen and rejuvenation cosmetics. Rejuvenation Res 21:91-101. <https://doi.org/10.1089/rej.2017.1931>

Kumar S, Singh A, Kumar B (2017) Identification and characterization of phenolics and terpenoids from ethanolic extracts of *Phyllanthus* species by HPLC-ESI-QTOF-MS/MS. J Pharm Anal 7:214-222. <https://doi.org/10.1016/j.jpha.2017.01.005>

Lange MK, Heberlé G, Milão D (2009) Avaliação da estabilidade e atividade antioxidante de uma emulsão base não-iônica contendo resveratrol. Braz J Pharm Sci 45:145-151. <https://doi.org/10.1590/S1984-82502009000100018>

Lee NYS, Khoo WKS, Adnan MA, Mahalingam TP, Fernandez AR, Jeevaratnam K (2016) The pharmacological potential of *Phyllanthus niruri*. J Pharm Pharmacol 68:953-969. <https://doi.org/10.1111/jphp.12565>

Leonardi GR, Gaspar LR, Campos PMBGM (2002) Estudo da variação do pH da pele humana exposta à formulação cosmética acrescida ou não das vitaminas A, E ou de ceramida, por metodologia não invasiva. An Bras Dermatol 77:563-569. <https://doi.org/10.1590/S0365-05962002000500006>

Mayba JN, Gooderham MJ (2018) A guide to topical vehicle formulations. J Cutan Med Surg 22(2):207-212.

Oliveira CA, Dario MF (2018) Bioactive Cosmetics. In: Martínez L, Kharissova O, Kharisov B (ed). Handbook of Ecomaterials. 1.ed. Springer International Publishing. 1-23.

Oliveira F (2014) Farmacognosia. 2.ed. São Paulo: Atheneu. 436p.

- Owabhel F, Eboh AS (2022) Antioxidant, chelating and HPLC quantification of phenols in extract of *Phyllanthus niruri*. WJPLS 3(1):1–6. <https://doi.org/10.53346/wjapls.2022.3.1.0029>
- Pianovski AR, Vilela AFG, Silva AAS da, Lima CG, Silva KK, Carvalho VFM, Musis CR, Machado SRP, Ferrari M (2008) Uso do óleo de pequi (*Caryocar brasiliense*) em emulsões cosméticas: desenvolvimento e avaliação da estabilidade física. Rev Bras Ciênc Farm 44:249–259. <https://doi.org/10.1590/S1516-93322008000200010>
- Rosário MS, Gauto MIR, Silva ACLN, Sales JS, Pereira FS, Santos EP (2021) Study of stability of cosmetic emulsion with potential of hydrating cream for the treatment of cutaneous xerosis using babassu oil (*Orbignya phalerata* Martius). Braz J Dev 7:29552–29570. <https://doi.org/10.34117/bjdv7n3-596>
- Shanbhag S, Nayak A, Narayan R, Nayak UY (2019) Anti-aging and sunscreens: paradigm shift in cosmetics. Adv Pharm Bull 9:348–359. <https://doi.org/10.15171/apb.2019.042>
- Soares CB, Soares RDF, Soares, GB, Ferreira EB, Ribeiro JC (2021) Development study of magistral cosmetic formulations containing 5% caffeine. Int J Dev Res 11(11):51753-51759.
- Souza VB, Ferreira JRN (2010) Desenvolvimento e estudos de estabilidade de cremes e géis contendo sementes e extratos do bagaço da uva Isabel (*Vitis labrusca* L.). Rev Ciênc Farm Básica Apl 31:217–222.